

Landau LITP/3-CP-93

PAR-LPTHE 93-34

June 1993

CRITICAL POINT CORRELATION FUNCTION FOR THE 2D RANDOM COUPLING ISING MODEL

A.L.Talapov*

*Landau Institute for Theoretical Physics**GSP-1 117940 Moscow V-334, Russia*

Vl.S.Dotsenko†

*LPTHE‡**Université Pierre et Marie Curie, PARIS VI**Tour 16, 1^{er} étage**4 place Jussieu**75252 Paris CEDEX 05, FRANCE**and**Landau Institute for Theoretical Physics**Moscow, Russia*

ABSTRACT

High accuracy Monte Carlo simulation results for 1024×1024 Ising system with ferromagnetic impurity bonds are presented. Spin-spin correlation function at a critical point is found to be numerically very close to that of a pure system. This is not trivial since a critical temperature for the system with impurities is almost two times lower than pure Ising T_c . Small deviations from the pure behaviour contradict to some of the competing theories.

*E-mail address: talapov@itp.sherna.msk.su

†E-mail address: dotsenko@lpthe.jussieu.fr

‡Laboratoire associé No. 280 au CNRS

Influence of impurities on a critical behaviour has been a subject of numerous papers.

For the simplest possible model – 2D Ising model this problem has been considered theoretically [1–10], experimentally [11, 12] and using computer simulations [13–18].

Most of the simulations were devoted to thermodynamic properties such as specific heat, magnetization and magnetic susceptibility. On the other hand, theories give direct predictions for spin-spin correlation function

$$\langle S(0)S(r) \rangle \quad (1)$$

where r is a distance between spins.

We used cluster algorithm special purpose processor (SPP) [19, 20] to get accurate values of $\langle S(0)S(r) \rangle$ at a critical point. The SPP realizes in hardware Wolff cluster algorithm, and therefore does not suffer of critical slowing down.

The SPP spends $375ns$ per one cluster spin. It also has a simple hardware for extremely fast calculation of spin-spin correlation functions. Time necessary to get the correlation function for some r is equal to $L^2 * 21ns$, where L is a linear lattice size.

We study the following model. Coupling constant J on each bond can take two positive values: J_1 with probability p and J_0 with probability $1 - p$. For $p = 0.5$ duality relation [21] shows that T_c is equal to that of a pure model with all horizontal bonds equal to J_1 and all vertical bonds equal to J_0 . This greatly simplifies simulation data analysis, and so we used $p = 0.5$.

Theoretical models use a small parameter

$$g \sim p(J_0 - J_1)^2 \quad (2)$$

This parameter is connected with impurity induced length l_i

$$\log l_i \sim \frac{1}{g} \quad (3)$$

To be able to notice deviations from the pure critical behaviour we should have

$$l_i \ll L \quad (4)$$

To satisfy this condition we used $L = 1024$ and quite different values of J : $J_0 = 1$ and $J_1 = 0.25$. So in the simulations g is not very small.

Theories deal with continuum limit and infinite lattice size. Simulations are conducted on a finite lattice with periodic boundary conditions. We calculated $\langle S(0)S(r) \rangle$ for spins, located along one lattice row. In this case distance r can take only integer values.

Our simulations for the pure case [20] showed that discrete lattice effects are significant for $r < 8$. Continuum theory can be applied for larger distances. But the finite size corrections for $r > 8$ are very significant and should be taken into account explicitly.

Pure Ising correlation function $c_0(r)$ for $r/L \rightarrow 0$ has been calculated in [22]

$$c_0(r) = \frac{0.70338}{r^{1/4}} \quad (5)$$

Continuum limit of (1) for the finite lattice with periodic boundary conditions $c(r, L)$ has been obtained in [23]

$$c(r, L) \sim \frac{\sum_{\nu=1}^4 \left| \theta_{\nu}\left(\frac{r}{2L}\right) \right|}{\left| \theta_1\left(\frac{r}{L}\right) \right|^{1/4}} \quad (6)$$

where θ are Jacobi theta functions. For our purposes $c(r, L)$ can be written in a simpler form

$$c(r, L) \approx A(L) \frac{1 + e^{-\pi/4} \left[\sin\left(\frac{\alpha}{2}\right) + \cos\left(\frac{\alpha}{2}\right) \right] + e^{-9\pi/4} \left[\cos\left(\frac{3\alpha}{2}\right) - \sin\left(\frac{3\alpha}{2}\right) \right]}{(\sin(\alpha) - e^{-2\pi} \sin(3\alpha))^{1/4}} \quad (7)$$

where $\alpha = \pi r/L$. The coefficient $A(L)$ can be obtained using expression for $c_0(r)$. Formula for $c(r, L)$ is in excellent agreement with simulation results [20] for the pure system.

In Fig.1 we show the ratio of computed $\langle S(0)S(r) \rangle$ to $c(r, L)$ for $L = 1024$.

To get mean values of $\langle S(0)S(r) \rangle$ and standard deviations we used one thousand samples with different impurities distribution. For each sample all spins initially were pointing in the same direction. Two thousand Wolff clusters were flipped to thermalize spin distribution at critical temperature. Another eight thousand clusters were flipped to calculate mean values of $\langle S(0)S(r) \rangle$ for each sample. For $L = 1024$ one cluster flip at T_c requires about 0.1sec.

Correlation functions for different r were measured for the same spin configurations.

Error bars are determined mainly by different behaviour of $\langle S(0)S(r) \rangle$ for different impurity distributions and not by thermal fluctuations for a given sample.

Deviations of $\langle S(0)S(r) \rangle$ from $c(r, L)$ at $r < 8$ are due to the discrete lattice effects. In fact, at these distances correlation function of the impure system is extremely close to the pure correlation function. For $r = 1$ difference between them is of the order of 10^{-3} . At first sight it seems to be quite natural, because $r \ll l_i$ and $\langle S(0)S(0) \rangle = 1$. But the critical temperatures for pure and impure cases are different, and continuum theory cannot exclude strong renormalization of $\langle S(0)S(1) \rangle$.

On the other hand, for $r > 8$ pure correlation function practically coincides with $c(r, L)$. So, Fig.1 shows that impurities decrease spin-spin correlations at large r . Again, this is not trivial because of the difference of critical temperatures for pure and impure cases.

This result contradicts the renormalization group calculation of DD [1] for the spin-spin correlation function, which would instead show increase if spin correlations, – see

also the discussion in the first paper of [13] where the numerical simulation problem has been defined.

The calculations of DD were based on the interacting fermion model [24, 25] representing the 2D Ising model with impurities [26]. The replica components trick has been used, for the fermion fields renormalization group calculation, which gave a new critical behavior for the correlation length and of the specific heat, as compared to the Ising model on a perfect lattice. The result of [26] has later been confirmed by calculations using other approaches, see [2–7], as well as by the computer simulations [13]. (See however [8,9] which claim a finite specific heat at T_c). In these calculations, of the energy operator related quantities, the replica components number N , to be taken eventually equal to 0, plays a minor role, that of the diagram counting, both in the original calculation in [26] using fermions, as well as in other approaches (bosonization, spin-component replicas)[2–7].

On the other hand, in the calculation of the spin–spin correlation function in [27] (see also [1]) the fermion replica components did play a dynamical role, while the other approaches, dealing more directly with the spin operators, employed replicas for diagram counting only.

Our present numerical result for the spin–spin function shows that fully dynamical effects due to replicas of the fermion fields lead to a wrong result, for this particular problem. The precise reason for this remains unclear.

The approaches based on bosonization gave the spin–spin function to be exactly same (in the scaling limit) as that of the pure Ising model. This is also not supported by our numerical result.

The deviations of $\langle SS \rangle$ in Fig.1 should be compared then with the standard renormalization group applied directly to the spin operators. The result of [2,5,6] was that the coefficient of the first order renormalization group equation vanishes, in the $N \rightarrow 0$ limit (N now is the spin operator replicas). No one had actually calculated the second order coefficient (let us denote it C_2), but assuming it is non–vanishing the result is standard. As the renormalization group equation one has:

$$\frac{d\Gamma(x)}{dx} = -C_2 g^2(x) \quad (8)$$

Here

$$\Gamma(x) = \frac{\langle S(0)S(r) \rangle}{\langle S(0)S(r) \rangle_0} \quad (9)$$

$$\langle S(0)S(r) \rangle_0 \sim c_0(r) \sim \frac{1}{r^{1/4}} \quad (10)$$

$$x = \log(r) \quad (11)$$

The renormalization group coupling constant $g(x)$ is [26]:

$$g(x) = \frac{g_0}{1 + \frac{4g_0}{\pi}x} \quad (12)$$

Then from (8) one finds:

$$\Gamma(x) = \exp\left(-\frac{C_2 g_0^2 x}{1 + \frac{4g_0}{\pi}x}\right) \quad (13)$$

and

$$\langle S(0)S(r) \rangle \sim \frac{1}{r^{1/4}} \exp\left(-\frac{C_2 g_0^2 x}{1 + \frac{4g_0}{\pi}x}\right) \quad (14)$$

So it is not exactly the pure Ising model behavior as there is a cross-over in the amplitude:

$$\begin{aligned} \Gamma(x) &\approx 1, & x &\ll x_i \\ &\approx \exp\left(-\frac{\pi C_2}{4}g_0\right), & x &\gg x_i \end{aligned} \quad (15)$$

$x_i = \pi/4g_0 = \log(l_i)$ is the impurities driven cross-over scale.

In the cross-over region $\langle SS \rangle / \langle SS \rangle_0 = \Gamma(x)$ is decreasing. This is what we observe in Fig.1.

Suppose that the finite lattice correlation function renormalization is given by (13) as well. Then we can estimate C_2 . Value of g_0 , found from the magnetic susceptibility, magnetization and specific heat data [13,28], is somewhere near 0.3. According to Fig.1 C_2 turns out to be quite small, of the order of 10^{-2} . This also agrees with the magnetic susceptibility and magnetization data [28].

More specifically, the estimate by the slope of the plot in Fig.1 for r relatively small, $8 < r < 32$, where the finite size corrections should still be insignificant, gives $C_2 \sim 0.03$. Clearly, this is an order of magnitude estimate. (Here, by finite size corrections, we mean in fact the small corrections to them due to the extra factor $\Gamma(x)$, eqs.(13),(14), in the correlation function of the impure model, as compared to the pure one).

We remark finally that previous measurements of the magnetization in [13] has not shown difference with the pure model because the precision was not sufficient. In fact, the variation we observe for $\langle SS \rangle / \langle SS \rangle_0$ corresponds, for small $4g_0/\pi x$, to an effective change in the magnetization index on the value $C_2 g_0^2/2$, which is from 0.125 to ~ 0.126 . This is just beyond the accuracy in [13].

Acknowledgements

We acknowledge useful discussions with V.B.Andreichenko, Vik.S.Dotsenko, V.L.Pokrovsky, W.Selke, and L.N.Shchur.

References

- [1] Vik.S.Dotsenko, Vl.S.Dotsenko, *Adv.Phys.* **32** (1983) 129
- [2] B.N.Shalaev, *Sov.Phys. Solid State* **26** (1984) 1811
- [3] R.Shankar, *Phys.Rev.Lett.* **58** (1987) 2466
- [4] R.Shankar, *Phys.Rev.Lett.* **61** (1988) 2390
- [5] A.B.Zamolodchikov, private communication (1987), unpublished
- [6] A.W.W.Ludwig, *Phys.Rev.Lett.* **61** (1988) 2388
- [7] A.W.W.Ludwig, *Nucl.Phys.* **B330** (1990) 639
- [8] P.N.Timonin, *Sov.Phys.JETP* **68** (1989) 512
- [9] K.Ziegler, *Nucl.Phys.* **B344** (1990) 557
- [10] I.O.Mayer, *J.Phys.* **A22** (1989) 2815
- [11] H.Ikeda, *J.Phys.Soc.Jpn.* **50** (1981) 3215
- [12] M.Hagen, R.A.Cowley, R.M.Nicklow, H.Ikeda, *Phys.Rev.* **B36** (1987) 401
- [13] V.B.Andreichenko, Vl.S.Dotsenko, W.Selke, J.-S.Wang, *Nucl.Phys.* **B344** (1990) 531
J.-S.Wang, W.Selke, Vl.S.Dotsenko, V.B.Andreichenko, *Europhys.Lett.*, **11** (1990) 301; *Physica* **A164** (1990) 221
- [14] A.L.Talapov, A.B.Andreichenko, Vl.S.Dotsenko and L.N.Shchur, *Sov.Phys.JETP Lett.* **51** (1990) 182. (*Pis'ma v Zh.Esp.Teor.Fiz* **51** (1990) 161.)
- [15] A.L.Talapov, A.B.Andreichenko, Vl.S.Dotsenko, and L.N.Shchur, in *Computer Simulations in Condensed Matter Physics III*, eds. D.P.Landau, K.K.Mon and H.B.Schuettler, (Springer, Heidelberg, 1991).
- [16] A.B.Andreichenko, Vl.S.Dotsenko, L.N.Shchur and A.L.Talapov, *Int.J.Mod.Phys.* **C2** (1991) 805.
- [17] M.Fahnle, T.Holey, J.Eckert ICM 91, Edinburgh to be published in PR B.
- [18] S.L.A.de Queiroz, R.B.Stinchcombe *Phys. Rev.* **B46** (1992) 6635
- [19] A.L.Talapov, L.N.Shchur, V.B.Andreichenko, Vl.S.Dotsenko, *Mod.Phys.Lett.* **B6** 1111

- [20] A.L.Talapov, V.B.Andreichenko, Vl.S.Dotsenko, L.N.Shchur, *Int.J.Mod.Phys.* **C** to appear
- [21] R.Fish *J.Stat.Phys.* **18** (1978) 111
- [22] T.T.Wu, B.M.McCoy, C.A.Tracy, E.Barouch, *Phys. Rev.* **B13** (1976) 31
- [23] P.di Francesco, H.Saleur, J.-B.Zuber, *Nucl. Phys.* **B290** [FS20] (1987) 527
- [24] P.K.Mitter and P.H.Weisz, *Phys. Rev.* **D8** (1973) 4410
- [25] D.J.Gross and A.Neveu, *Phys.Rev.* **D10** (1974) 3235
- [26] Vik.S.Dotsenko and Vl.S.Dotsenko, *Sov.Phys. JETP Lett.* **33** (1981) 37
- [27] Vik.S.Dotsenko and Vl.S.Dotsenko, *J.Phys.* **C15** L557 (1982)
- [28] A.L.Talapov, A.B.Andreichenko, Vl.S.Dotsenko and L.N. Shchur, to be published

Figure Captions

Fig.1. Ratio of $\langle S(0)S(r) \rangle$ to $c(r, L)$. Solid line connects data points for the system with impurities at $(1/T_c) = .8070519$. Dashed line shows data for the pure Ising model [23] at $(1/T_c) = .4406868$.